

# Seroprevalence of antibodies against bacterial pathogens in sheep from Equatorial Guinea

D. Loureiro <sup>(1)</sup>, L.F. Moura-Costa <sup>(1)</sup>, R.S. Jordão <sup>(2)</sup>, N.C. Menezes <sup>(3)</sup>,  
E.S. Macedo <sup>(4)</sup>, T.S. Guimarães <sup>(2)</sup>, R. Meyer <sup>(1)</sup> & R.W. Portela <sup>(1)\*</sup>

(1) Laboratório de Imunologia e Biologia Molecular, Instituto de Ciências da Saúde, Universidade Federal da Bahia, Av. Reitor Miguel Calmon s/n, Vale do Canela, Salvador, BA 40110-100, Brazil

(2) Laboratório de Produção de Imunobiológicos, Instituto Biológico de São Paulo, São Paulo, SP 04014-002, Brazil

(3) Associação de Criadores de Caprinos e Ovinos da Bahia, Salvador, BA 41611-670, Brazil

(4) ARG Ltda. – Guinée Équatoriale, Oyala, Wele Nzaz, Equatorial Guinea

\*Corresponding author: rwportela@gmail.com

## Summary

Equatorial Guinea (EG) is a country in Central Africa with typical tropical weather. Sheep are an important source of food in EG, but the absence of information regarding infectious diseases that affect the native flocks of small ruminants is a concern. The country is currently implementing several new practices in the sheep industry associated with the importation of sheep from other countries. This study aimed to evaluate the seroprevalence of bacterial infections that are important to the sheep industry in EG sheep flocks. The detection of specific antibodies for the four agents studied was performed using enzyme-linked immunosorbent assay in 1,002 serum samples from EG sheep. The results showed a true prevalence of 13.37% for antibodies against *Corynebacterium pseudotuberculosis*, 0.59% for *Brucella ovis*, 19.89% for *Chlamydophila abortus* and 0.79% for *Mycoplasma agalactiae* in animals from production flocks. Among a group of 35 samples from isolated native animals, 47.56% were seropositive for antibodies against *C. pseudotuberculosis*, 42.84% for *B. ovis*, 54.28% for *C. abortus* and 11.35% for *M. agalactiae*. These results comprise the first report of the prevalence of infectious diseases in sheep in EG. They highlight the importance of adopting adequate measures to prevent infection by bacteria endemic to EG native flocks during the development of the sheep industry in the country.

## Keywords

Caseous lymphadenitis – Contagious agalactia – Enzootic abortion – Equatorial Guinea – Ovine brucellosis – Sheep.

## Introduction

The Republic of Equatorial Guinea (EG) is located in West Africa (3°21'0"N latitude and 8°40'0"E longitude). It has a typical tropical climate, hot and rainy (26°C and 2,500 mm rain per year, on average) with high air humidity (80–90%). According to the Food and Agriculture Organization of the United Nations (FAO) (1), the estimated sheep population of EG comprises 5,000 animals, which are usually maintained in an extensive system. However, there are no published scientific studies regarding livestock diseases in the country.

Caseous lymphadenitis (CLA), caused by *Corynebacterium pseudotuberculosis*, is highly prevalent in many countries worldwide and causes decreases in weight, lactation and commercial value of the fleece and, in some cases, infertility

(2). In sheep, *Brucella ovis* infection leads to reduced fertility, management problems in the breeding season, abortion, epididymitis, orchitis and high rates of mortality among newborn lambs (3). Enzootic abortion of goats and sheep (EAGS), caused by the bacterium *Chlamydophila abortus*, is a zoonosis and a major cause of abortion, stillbirth, retained placenta, metritis, persistent cough, polyarthritis and keratoconjunctivitis (4), as well as a factor for consideration during animal transit (5). *Mycoplasma agalactiae* is the aetiological agent of contagious agalactia of goats and sheep (CAGS), a disease that causes high levels of morbidity and mortality; it is associated with mastitis, agalactia, arthritis, keratitis, reduced milk production and transient fever (6).

The economic losses caused by these four diseases are considerable (7, 8). Given the lack of information on infectious diseases in EG, and the fact that the country is

currently investing in the sheep industry, the aim of this study was to evaluate the prevalence of these four infections in EG sheep flocks.


## Materials and methods

Based on FAO estimates (1), the minimum sample size 'n' was calculated as proposed in (9), with a maximum sampling error of 5% and a 95% confidence level. With the aim of understanding the spread of the four diseases in the growing EG flock, samples were collected from several cities belonging to three of the four continental provinces, as shown in Figure 1. There are no data on sheep breeding in the insular provinces of EG.

Serum samples were collected from 1,002 sheep between 2013 and 2014, with the producers' consent. These serum samples were divided into two major groups: 967 sheep from properties that were investing in breeding enhancements, where sheep are raised in a semi-intensive management system, composed the first group; the animals that belonged to these properties and were included in this study are referred to as 'production animals'. Thirty-five native geographically isolated sheep composed the other

group; there were no reports of contact of these animals with other production animals, except wildlife. As these animals constituted a distinct group that may represent the infectivity profile of sheep in EG before the importation of other animals, these sheep are referred to as 'native animals'. Blood samples were collected in vacuum tubes, centrifuged at  $1,500 \times g$  for 10 min to obtain serum, distributed in 1.5 ml aliquots and stored in a freezer at  $-20^{\circ}\text{C}$ .

Serum samples were tested using an indirect enzyme-linked immunosorbent assay (ELISA) with 99% specificity and 89% sensitivity, as described in (10), with the objective of identifying the presence of immunoglobulin (Ig)G antibodies against *C. pseudotuberculosis*. Briefly, polystyrene microplates (Corning Life Sciences, Lowell, United States of America [USA]) were sensitised with *C. pseudotuberculosis* secreted/excreted antigen diluted 1:100 in carbonate-bicarbonate buffer (pH 9.6) and incubated at  $4^{\circ}\text{C}$  overnight. The plates were then washed twice with phosphate-buffered solution (PBS) 0.05% Tween 20 (PBST), blocked with 5% skimmed milk in PBST, and incubated at  $37^{\circ}\text{C}$  for 2 h. After two additional washes with PBST, 100  $\mu\text{l}$  of serum diluted 1:100 in PBST with 1% skimmed milk was added and the plates were incubated at  $37^{\circ}\text{C}$  for 1 h. Each serum sample was tested in duplicate. The plates were washed four times



**Fig. 1**  
**Map of Equatorial Guinea showing the division of the country into seven provinces, three in the insular region (Annobón, Bioco Norte and Bioco Sur) and four in the continental region (Litoral, Centro Sur, Wele-Nzas and Kié-Ntem)**


Cities in which the study's ovine serum samples were collected are represented by black dots. The numbers beside each black dot represent how many production flock animals were sampled at each location, and the numbers followed by the symbol '\*' represent the quantity of isolated native animals that were sampled at two different locations

with PBST and a rabbit anti-ovine IgG antibody conjugated to horseradish peroxidase (Serotec, Raleigh, USA), diluted 1:20,000, was added and incubated at 37°C for 45 min. After incubation, the plates were washed and the reaction was developed with 100 µl/well of a solution containing hydrogen peroxide and tetramethylbenzidine (Moss Inc., Pasadena, USA) for 15 min, and stopped with 4 N sulphuric acid. The results were read using an ELISA plate reader (Bio-Rad Laboratories, Hercules, USA) at 450 nm.

To identify the presence in sera of specific IgG antibodies against *B. ovis*, *C. abortus* and *M. agalactiae*, commercial indirect ELISA kits were used (Idexx Laboratories Inc., Westbrook, USA), all of which had 99.9% sensitivity and specificity, as reported by the manufacturer. The assays were performed following the manufacturer's protocol. All sera that tested positive for *B. ovis* antibodies had their status confirmed by agar gel immunodiffusion (Tecpar-PR, Curitiba, Brazil), following the protocol described by the manufacturer, with 96.4% sensitivity and 100% specificity (11). The true prevalence rates, adjusted for the sensitivity and specificity of each assay, were obtained by following the methodology described in (12).

## Results and discussion

The results are presented in Figure 2. An overall true seroprevalence of 19.31%, 19.83%, 0.65% and 0.83% for *C. pseudotuberculosis*, *C. abortus*, *B. ovis* and *M. agalactiae*, respectively, was found in sheep in EG. In production animals ( $n = 967$ ), true prevalence values of 13.37%,



**Fig. 2**  
**True prevalence of specific antibodies against bacterial pathogens in isolated native sheep and production flock animals from Equatorial Guinea**

The prevalence values were adjusted as proposed in (12), and the true prevalence estimates are shown at the top of the bars as percentages (%)

19.89%, 0.59% and 0.79% were estimated, while the isolated native animals ( $n = 35$ ) showed true prevalence values of 47.56%, 54.28%, 42.84% and 11.35% for specific antibodies against *C. pseudotuberculosis*, *C. abortus*, *B. ovis* and *M. agalactiae*, respectively.

The traditional extensive management system, which is commonly used in underdeveloped and developing countries, contributes not only to low productivity in the flocks, but also to the maintenance of many diseases (13). Moreover, the subclinical characteristics of the diseases studied here, and the lack of scientific information and infection reports in many of these countries, are factors that must be taken into consideration when establishing government programmes for the development and extension of the sheep industry.

The high seroprevalence of *C. pseudotuberculosis* and *C. abortus* in the native animals indicates the endemic presence of these bacteria in EG. This may be explained by the extensive management system, the absence of proper hygiene and the presence of risk factors associated with these infections in EG (14). There was also a significant prevalence of these infections in the animals in production flocks. According to (4), EAGS has a significant prevalence in all regions where goats and sheep are bred, which was confirmed in this work.

The results of this study showed that a high proportion of the isolated native animals were seropositive for *B. ovis*, but this was not replicated in the production animals. The low prevalence of the disease in production animals can be explained by the fact that the countries of origin of these animals (many were imported to EG as part of a sheep industry implementation programme) perform rigorous brucellosis control measures.

The low seroprevalence of *M. agalactiae* found in both groups in this study conflicts with previous research in other countries (6); this may be due to the low rates of animal transport and commercialisation in EG, and the low prevalence in native animals that are isolated from other sources of infection. Furthermore, because the production sheep may have been exposed to good hygiene conditions in their original countries, the disease appeared to be controlled in these animals.

The population of isolated native animals studied was small when compared with the total sample size (35 of 1,002 animals). This was because these native animals were the only ones identified that had never had any contact with imported animals, and therefore reflected the situation of the country before the development of a broader sheep industry. In these native animals, the prevalence of *B. ovis* and *M. agalactiae* infection was much greater than that found in production flocks. This represents a serious risk for the sheep industry implementation and extension programme

that is currently being conducted in EG, as has already been described in other countries (13). Interestingly, the initial concern was the reverse: that imported animals could act as carriers of diseases that were not present in native EG flocks.

## Conclusions

Sheep from EG present high levels of infection with *C. pseudotuberculosis* and *C. abortus*, with seroprevalence values of 47.56% and 54.28%, respectively. The prevalence of specific antibodies against *M. agalactiae* and *B. ovis* in isolated native animals reveals that the diseases caused

by these bacterial pathogens must be considered when implementing hygiene and preventive management procedures for animals that are being imported into the country.

## Acknowledgements

The authors are grateful to Francisca Soares (Laboratório de Imunologia e Biologia Molecular, Universidade Federal da Bahia, Salvador, Brazil) for technical assistance.

# Séroprévalence d'anticorps dirigés contre des agents pathogènes bactériens chez les moutons de Guinée équatoriale

D. Loureiro, L.F. Moura-Costa, R.S. Jordão, N.C. Menezes, E.S. Macedo, T.S. Guimarães, R. Meyer & R.W. Portela

### Résumé

La Guinée équatoriale est un pays d'Afrique centrale au climat typiquement tropical. Les moutons constituent une source importante d'aliments pour le pays, ce qui rend particulièrement alarmante l'absence d'informations concernant les maladies infectieuses affectant le cheptel autochtone de petits ruminants. Le pays met actuellement en place de nouvelles pratiques d'élevage ovin, en lien avec l'importation de moutons d'autres pays. La présente étude avait pour objet d'évaluer la prévalence sérologique de quelques infections bactériennes majeures pour le secteur de l'élevage dans le cheptel ovin de la Guinée équatoriale. Pour les besoins de l'étude, 1 002 échantillons de sérum prélevés de moutons autochtones ont été soumis à une épreuve immuno-enzymatique pour la détection d'anticorps spécifiques dirigés contre quatre agents bactériens. La prévalence réelle de moutons issus de troupeaux d'élevage possédant des anticorps vis-à-vis de chacune des bactéries recherchées s'est élevée à 13,37 % pour *Corynebacterium pseudotuberculosis*, à 0,59 % pour *Brucella ovis*, à 19,89 % pour *Chlamydomphila abortus* et à 0,79 % pour *Mycoplasma agalactiae*. Dans un groupe de 35 prélèvements issus d'animaux isolés autochtones, le taux d'échantillons positifs était de 47,56 % pour *C. pseudotuberculosis*, de 42,84 % pour *B. ovis*, de 54,28 % pour *C. abortus* et de 11,35 % pour *M. agalactiae*. Ces résultats constituent le premier rapport sur la prévalence des maladies infectieuses chez le mouton en Guinée équatoriale. Ils soulignent l'importance d'adopter des mesures appropriées afin de prévenir les infections dues à des bactéries endémiques dans le cheptel autochtone de la Guinée équatoriale, à un moment où ce pays cherche à développer l'élevage ovin.

### Mots-clés

Agalaxie contagieuse – Avortement enzootique – Brucellose ovine – Guinée équatoriale – Lymphadénite caséuse – Mouton.

# Seroprevalencia de anticuerpos contra patógenos bacterianos en ovejas de Guinea Ecuatorial

D. Loureiro, L.F. Moura-Costa, R.S. Jordão, N.C. Menezes, E.S. Macedo, T.S. Guimarães, R. Meyer & R.W. Portela

## Resumen

Guinea Ecuatorial es un país centroafricano con un clima tropical típico, en el que las ovejas constituyen una importante fuente de alimento. Por ello es preocupante la ausencia de información sobre las enfermedades infecciosas que afectan a los rebaños autóctonos de pequeños rumiantes. El país está implantando ahora una serie de nuevos procedimientos en el sector ovino, vinculados a la importación de ovejas de otros países. Los autores describen un estudio encaminado a evaluar la seroprevalencia en los rebaños de ovejas del país de infecciones bacterianas de importancia para el sector. Para detectar anticuerpos específicos contra los cuatro agentes estudiados se aplicó la técnica del ensayo inmunoenzimático a 1002 muestras de suero de ovejas autóctonas. Los resultados pusieron de manifiesto una prevalencia real de anticuerpos contra *Corynebacterium pseudotuberculosis* del 13,37%, del 0,59% contra *Brucella ovis*, del 19,89% contra *Chlamydophila abortus* y del 0,79% contra *Mycoplasma agalactiae* en ejemplares pertenecientes a rebaños productivos. De un grupo de 35 muestras procedentes de animales autóctonos aislados, un 47,56% resultaron seropositivas para los anticuerpos contra *C. pseudotuberculosis*, un 42,84% en el caso de *B. ovis*, un 54,28% en el de *C. abortus* y un 11,35% en el de *M. agalactiae*. Estos resultados, que constituyen el primer informe sobre la prevalencia de enfermedades infecciosas en las ovejas de Guinea Ecuatorial, demuestran la importancia de instituir medidas adecuadas para prevenir la infección por bacterias endémicas de los rebaños autóctonos de Guinea Ecuatorial en un momento en que la industria ovina del país está en pleno crecimiento.

## Palabras clave

Aborto enzoótico – Agalaxia contagiosa – Brucelosis ovina – Guinea Ecuatorial – Oveja – Seudotuberculosis de los ovinos.



## References

1. Food and Agriculture Organization of the United Nations (FAO) (2014). – Statistics Division. Available at: <http://faostat.fao.org> (accessed on 8 February 2017).
2. Dorella FA., Pacheco L.G.C., Seyffert N., Portela R.W., Meyer R., Miyoshi A. & Azevedo V. (2009). – Antigenes of *Corynebacterium pseudotuberculosis* and prospects for vaccine development. *Expert Rev. Vaccines*, **8** (2), 205–213. doi:10.1586/14760584.8.2.205.
3. Ridler A.L. & West D.M. (2011). – Control of *Brucella ovis* infection in sheep. *Vet. Clin. North Am. Food Anim. Pract.*, **27** (1), 61–66. doi:10.1016/j.cvfa.2010.10.013.
4. World Organisation for Animal Health (OIE) (2016). – Infection with *Chlamydophila abortus* (enzootic abortion of ewes, ovine chlamydiosis), Chapter 14.4. In *Terrestrial Animal Health Code, Vol. II. Recommendations applicable to OIE Listed diseases and other diseases of importance to international trade*. OIE, Paris. Available at: [www.oie.int/fileadmin/Home/eng/Health\\_standards/tahc/current/chapitre\\_chlamydophila\\_abortus.pdf](http://www.oie.int/fileadmin/Home/eng/Health_standards/tahc/current/chapitre_chlamydophila_abortus.pdf) (accessed on 8 February 2017).
5. Entrican G., Wheelhouse N., Wattedegera S.R. & Longbottom D. (2012). – New challenges for vaccination to prevent chlamydial abortion in sheep. *Comp. Immunol. Microbiol. Infect. Dis.*, **35** (3), 271–276. doi:10.1016/j.cimid.2011.12.001.

6. Gómez-Martín A., Amores J., Paterna A. & De la Fe C. (2013). – Contagious agalactia due to *Mycoplasma* spp. in small dairy ruminants: epidemiology and prospects for diagnosis and control. *Vet. J.*, **198** (1), 48–56. doi:10.1016/j.tvjl.2013.04.015.
  7. Guimarães A.S., Carmo F.B., Heinemann M.B., Portela R.W., Meyer R., Lage A.P., Seyffert N., Miyoshi A., Azevedo V. & Gouveia A.M. (2011). – High sero-prevalence of caseous lymphadenitis identified in slaughterhouse samples as a consequence of deficiencies in sheep farm management in the state of Minas Gerais, Brazil. *BMC Vet. Res.*, **7** (1), 68–76. doi:10.1186/1746-6148-7-68.
  8. Muma J.B., Samui K.L., Siamudaala V.M., Oloya J., Matop G., Omer M.K., Munyeme M., Mubita C. & Skjerve E. (2006). – Prevalence of antibodies to *Brucella* spp. and individual risk factors of infection in traditional cattle, goats and sheep reared in livestock–wildlife interface areas of Zambia. *Trop. Anim. Hlth Prod.*, **38** (1), 195–206. doi:10.1007/s11250-006-4320-9.
  9. Sreenivasaiah P.V. (2008). – Veterinary biostatistics, 1st Ed. International Book Distributing Company, Lucknow, India, 487 pp.
  10. Rebouças M.F., Loureiro D., Bastos B.L., Moura-Costa L.F., Hanna S.A., Azevedo V., Meyer R. & Portela R.W. (2013). – Development of an indirect ELISA to detect *Corynebacterium pseudotuberculosis* specific antibodies in sheep employing T1 strain culture supernatant as antigen. *Braz. J. Vet. Res.*, **33** (1), 1296–1302. doi:10.1590/s0100-736x2013001100002.
  11. Marín C.M., Bagués M.P.J., Blasco J.M., Gamazo C., Moriyón I. & Díaz R. (1989). – Comparison of three serological tests for *Brucella ovis* infection of rams using different antigenic extracts. *Vet. Rec.*, **125** (1), 504–508. doi:10.1136/vr.125.20.504.
  12. Noordhuizen J.P.T.M., Thrusfield M.V., Frankena K. & Graat E.A.M. (2001). – Application of quantitative methods in veterinary epidemiology, 1st Ed. Wageningen Academic Publishers, Wageningen, the Netherlands, 429 pp. doi:10.3920/978-90-74134-89-7.
  13. Sherman D.M. (2011). – The spread of pathogens through trade in small ruminants and their products. In The spread of pathogens through international trade (S.C. MacDiarmid, ed.). *Rev. Sci. Tech. Off. Int. Epiz.*, **30** (1), 207–217. doi:10.20506/rst.30.1.2036.
  14. Kaba J., Nowicki M., Frymus T., Nowicka D., Witkowski L., Szaluś-Jordanow O., Czopowicz M. & Thrusfield M. (2011). – Evaluation of the risk factors influencing the spread of caseous lymphadenitis in goat herds. *Pol. J. Vet. Sci.*, **14** (1), 231–237. doi:10.2478/v10181-011-0035-6.
-